

Table S1. Total species list (alphabetical by English common name) and proportion of three forests occupied for all species observed, calculated as naïve occupancy estimates.

| English name | Scientific name | Manual Restoration | Natural Regrowth | Shifting Cultivation |
|---------------------------|---------------------------------|---------------------------|-------------------------|-----------------------------|
| Andean solitaire | <i>Myadestes ralloides</i> | 0.0 | 0.0 | 0.2 |
| Bananaquit | <i>Coereba flaveola</i> | 0.8 | 1.0 | 0.8 |
| Bay-headed tanager | <i>Tangara gyrola</i> | 0.6 | 1.0 | 1.0 |
| Black and white becard | <i>Pachyramphus albogriseus</i> | 0.2 | 0.0 | 0.0 |
| Black caracara | <i>Daptrius ater</i> | 0.0 | 1.0 | 0.2 |
| Black-billed thrush | <i>Turdus Ignobilis</i> | 0.2 | 1.0 | 0 |
| Black-crowed tityra | <i>Tityra inquisitor</i> | 0.0 | 0.5 | 0 |
| Black-faced dacnis | <i>Dacnis lineata</i> | 0.2 | 1.0 | 0.6 |
| Blackpoll warbler | <i>Dendroica striata</i> | 0.4 | 0 | 0 |
| Black-tailed tityra | <i>Tityra cayana</i> | 0.2 | 0.5 | 0.2 |
| Blue dacnis | <i>Dacnis cayana</i> | 0.0 | 0.5 | 0.2 |
| Blue-crowed trogon | <i>Trogon curucui</i> | 0.0 | 0 | 0.6 |
| Blue-fronted lancebill | <i>Doryfera johannae</i> | 0.2 | 0.0 | 0.2 |
| Blue-gray tanager | <i>Thraupis episcopus</i> | 1.0 | 1.0 | 0.4 |
| Blue-headed parrot | <i>Pionus menstruus</i> | 0.2 | 0.5 | 0.2 |
| Blue-necked tanager | <i>Tangara cyanicollis</i> | 1.0 | 1.0 | 1.0 |
| Boat-billed flycatcher | <i>Megarynchus pitangua</i> | 1.0 | 1.0 | 0.6 |
| Broad-winged hawk | <i>Buteo platypterus</i> | 0.2 | 0.0 | 0.0 |
| Buff-throated saltator | <i>Saltator maximus</i> | 0.2 | 0.5 | 0.0 |
| Buff-throated woodcreeper | <i>Xiphorhynchus guttatus</i> | 0.8 | 0.5 | 0.4 |
| Canada warbler | <i>Wilsonia canadensis</i> | 0.0 | 0.5 | 0.2 |
| Cattle egret | <i>Bubulcus ibis</i> | 0.0 | 1.0 | 0.0 |

| | | | | |
|--------------------------------|----------------------------------|-----|-----|-----|
| Chestnut-breasted chlorophonia | <i>Chlorophonia pyrrhophrys</i> | 0.0 | 0.5 | 0.0 |
| Chestnut-capped puffbird | <i>Bucco macrodactylus</i> | 0.0 | 0.0 | 0.2 |
| Chestnut-crowned becard | <i>Pachyramphus castaneus</i> | 0.0 | 0.0 | 0.2 |
| Dusky-capped flycatcher | <i>Myiarchus tuberculifer</i> | 1.0 | 1.0 | 0.2 |
| Eastern wood pewee | <i>Contopus virens</i> | 0.0 | 1.0 | 0.0 |
| Fawn-breasted tanager | <i>Pipraeidea melanonota</i> | 0.0 | 0.5 | 0.0 |
| Fiery-throated fruiteater | <i>Pipreola chlorolepidota</i> | 0.0 | 0.0 | 0.2 |
| Forest-elaenia | <i>Myiopagis gaimardii</i> | 0.0 | 0.0 | 0.4 |
| Fork-tailed woodnymph | <i>Thalurania furcata</i> | 0.2 | 0.5 | 0.4 |
| Fulvous-breasted flatbill | <i>Rhynchocyclus fulvipectus</i> | 0.0 | 0.5 | 0.0 |
| Gilded barbet | <i>Capito auratus</i> | 0.0 | 1.0 | 0.2 |
| Glittering-throated emerald | <i>Amazilia fimbriata</i> | 0.6 | 1.0 | 0.2 |
| Golden tanager | <i>Tangara arthus</i> | 0.0 | 0.0 | 0.2 |
| Golden-olive woodpecker | <i>Colaptes rubiginosus</i> | 0.0 | 0.0 | 0.2 |
| Golden-tailed sapphire | <i>Chrysuronia oenone</i> | 0.4 | 0.0 | 0.2 |
| Gray-chinned hermit | <i>Phaethornis griseogularis</i> | 0.6 | 0.5 | 0.2 |
| Grayish saltator | <i>Saltator coerulescens</i> | 1.0 | 0.5 | 0.6 |
| Great jacamar | <i>Jacamerops aureus</i> | 0.0 | 0.0 | 0.2 |
| Great kiskadee | <i>Pitangus sulphuratus</i> | 0.8 | 0.0 | 0.0 |
| Great-billed hermit | <i>Phaethornis malaris</i> | 1.0 | 0.5 | 0.4 |
| Green-and-gold tanager | <i>Tangara schrankii</i> | 1.0 | 1.0 | 1.0 |
| Lafresnaye's piculet | <i>Picumnus lafresnayi</i> | 0.6 | 0.0 | 0.2 |
| Lineated woodpecker | <i>Dryocopus lineatus</i> | 0.4 | 0.0 | 0.2 |

| | | | | |
|-----------------------------|--------------------------------------|-----|-----|-----|
| Lined antshrike | <i>Thamnophilus zarumae</i> | 0.2 | 0.0 | 0.0 |
| Long-tailed tyrant | <i>Colonia colonus</i> | 0.0 | 0.0 | 0.4 |
| Magpie tanager | <i>Cissopis leveriana</i> | 0.8 | 1.0 | 0.6 |
| Masked tityra | <i>Tityra semifasciata</i> | 0.2 | 0.0 | 0.2 |
| Olivaceous siskin | <i>Carduelis olivacea</i> | 0.4 | 0.0 | 0.2 |
| Olivaceous woodcreeper | <i>Sittasomus griseicapillus</i> | 0.8 | 1.0 | 1.0 |
| Olive sided flycatcher | <i>Contopus cooperi</i> | 0.4 | 0.0 | 0.2 |
| Orange-bellied euphonia | <i>Euphonia xanthogaster</i> | 0.6 | 1.0 | 1.0 |
| Palm tanager | <i>Thraupis palmarum</i> | 1.0 | 1.0 | 1.0 |
| Paradise tanager | <i>Tangara chilensis</i> | 1.0 | 1.0 | 1.0 |
| Pauraque | <i>Nyctidromus albicollis</i> | 0.0 | 0.5 | 0.0 |
| Plain-brown woodcreeper | <i>Dendrocincla fuliginosa</i> | 0.2 | 0.0 | 0.0 |
| Red-headed barbet | <i>Eubucco bourcierii</i> | 0.0 | 1.0 | 0.0 |
| Ringed kingfisher | <i>Megaceryle torquata</i> | 0.0 | 0.5 | 0.0 |
| Roadside hawk | <i>Buteo magirostris</i> | 0.0 | 0.5 | 0.4 |
| Ruddy pigeon | <i>Patagioenas subvinacea</i> | 0.0 | 0.5 | 0.2 |
| Rufous-collared sparrow | <i>Zonotrichia capensis</i> | 0.0 | 0.0 | 0.2 |
| Russet-backed oropendola | <i>Psarocolius angustifrons</i> | 1.0 | 1.0 | 1.0 |
| Saffron-crowned tanager | <i>Tangara xanthocephala</i> | 0.2 | 0.0 | 0.0 |
| Scarlet tanager | <i>Piranga olivacea</i> | 0.0 | 1.0 | 0.0 |
| Silver-beaked tanager | <i>Ramphocelus carbo</i> | 1.0 | 1.0 | 0.6 |
| Smoke-colored pewee | <i>Contopus fumigatus</i> | 0.2 | 0.0 | 0.0 |
| Smokey-brown woodpecker | <i>Veniliornis fumigatus</i> | 0.2 | 0.0 | 0.6 |

| | | | | |
|-------------------------------|----------------------------------|-----|-----|-----|
| Social flycatcher | <i>Myiozetetes similis</i> | 0.4 | 1.0 | 1.0 |
| Southern rough-winged swallow | <i>Stelgidopteryx ruficollis</i> | 0.2 | 0.0 | 0.0 |
| Speckled chachalaca | <i>Ortalis guttata</i> | 0.2 | 0.0 | 0.0 |
| Spot-breasted woodpecker | <i>Colaptes punctigula</i> | 0.0 | 0.5 | 0.2 |
| Spotted tanager | <i>Tangara punctata</i> | 0.4 | 1.0 | 0.4 |
| Squirrel cuckoo | <i>Piaya cayana</i> | 0.4 | 0.0 | 0.4 |
| Streaked xenops | <i>Xenops rutilans</i> | 0.0 | 0.0 | 0.2 |
| Swainson' s thrush | <i>Catharus ustulatus</i> | 0.4 | 0.5 | 0.4 |
| Swallow tanager | <i>Tersina viridis</i> | 0.2 | 0.5 | 0.0 |
| Tropical kingbird | <i>Tyrannus melancholicus</i> | 0.4 | 0.5 | 0.2 |
| Turkey vulture | <i>Cathartes aura</i> | 0.0 | 0.0 | 0.2 |
| Turquoise tanager | <i>Tangara mexicana</i> | 0.0 | 0.5 | 0.0 |
| Vermilion tanager | <i>Calochaetes coccineus</i> | 0.0 | 0.5 | 0.0 |
| Violaceous jay | <i>Cyanocorax violaceus</i> | 1.0 | 1.0 | 0.8 |
| Violet-fronted brilliant | <i>Heliodoxa leadbeateri</i> | 0.0 | 0.0 | 0.4 |
| Warbling antbird | <i>Hypocnemis peruviana</i> | 0.0 | 0.0 | 0.2 |
| Wedge-billed woodcreeper | <i>Glyphorynchus spirurus</i> | 0.4 | 0.0 | 0.4 |
| White-collared swift | <i>Streptoprocne zonaris</i> | 0.2 | 0.5 | 0.4 |
| White-crested elaenia | <i>Elaenia albiceps</i> | 0.2 | 0.0 | 0.0 |
| White-lined tanager | <i>Tachyphonus rufus</i> | 0.2 | 0.0 | 0.2 |
| White-necked jacobin | <i>Florisuga mellivora</i> | 0.0 | 0.0 | 0.2 |
| White-throated toucan | <i>Ramphastos tucanus</i> | 0.0 | 0.0 | 0.4 |
| Yellow browed sparrow | <i>Ammodramus aurifrons</i> | 0.4 | 0.5 | 0.0 |
| Yellow-backed tanager | <i>Hemithraupis flavicollis</i> | 0.0 | 0.0 | 0.2 |

| | | | | |
|---------------------------------|---------------------------------------|-----|-----|-----|
| Yellow-rumped cacique | <i>Cacicus cela</i> | 1.0 | 1.0 | 0.2 |
| Yellow-throated bush tanager | <i>Chlorospingus flavigularis</i> | 0.0 | 0.0 | 0.2 |
| Yellow-tufted woodpecker | <i>Melanerpes cruentatus</i> | 0.2 | 0.5 | 0.4 |

Table S2. Models describing forest occupancy for 18 bird species, ranked by (Q)AICc, with $\Delta(Q)AICc$ and model weights.

a. Bananaquit

| Model | AIC | ΔAIC | Model weight |
|--|------------|--------------------------------|---------------------|
| $\Psi(.) p(.)$ | 141.90 | 0.00 | 0.6735 |
| $\Psi(.) p(\text{Forest})$ | 144.26 | 2.36 | 0.2069 |
| $\Psi(\text{Forest}) p(.)$ | 145.89 | 3.99 | 0.0916 |
| $\Psi(\text{Forest}) p(\text{Forest})$ | 148.26 | 6.36 | 0.0280 |

b. Bay-headed tanager

| Model | AIC | ΔAIC | Model weight |
|--|------------|--------------------------------|---------------------|
| $\Psi(.) p(\text{Forest})$ | 157.76 | 0.00 | 0.8248 |
| $\Psi(\text{Forest}) p(\text{Forest})$ | 161.62 | 3.86 | 0.1197 |
| $\Psi(.) p(.)$ | 164.16 | 6.40 | 0.0336 |
| $\Psi(\text{Forest}) p(.)$ | 165.02 | 7.26 | 0.0219 |

c. Black vulture

| Model | QAIC | $\Delta QAIC$ | Model weight |
|--|-------------|---------------------------------|---------------------|
| $\Psi(\text{Forest}) p(.)$ | 70.86 | (-)0 | 0.4577 |
| $\Psi(.) p(\text{Forest})$ | 70.86 | (-)0 | 0.4577 |
| $\Psi(\text{Forest}) p(\text{Forest})$ | 74.86 | 4.00 | 0.0619 |
| $\Psi(.) p(.)$ | 76.87 | 6.01 | 0.0227 |

d. Black-faced dacnis

| Model | QAIC | $\Delta QAIC$ | Model weight |
|--|-------------|---------------------------------|---------------------|
| $\Psi(.) p(\text{Forest})$ | 44.06 | 0.00 | 0.6281 |
| $\Psi(.) p(.)$ | 46.16 | 2.10 | 0.2198 |
| $\Psi(\text{Forest}) p(\text{Forest})$ | 48.00 | 3.94 | 0.0876 |
| $\Psi(\text{Forest}) p(.)$ | 48.61 | 4.55 | 0.0646 |

e. Blue-necked tanager

| Model | AIC | ΔAIC | Model weight |
|--|------------|--------------------------------|---------------------|
| $\Psi(.) p(\text{Forest})$ | 221.78 | 0.00 | 0.8808 |
| $\Psi(\text{Forest}) p(\text{Forest})$ | 225.78 | 4.00 | 0.1192 |
| $\Psi(.) p(.)$ | 262.09 | 40.31 | 0.0000 |
| $\Psi(\text{Forest}) p(.)$ | 264.24 | 42.46 | 0.0000 |

f. Blue-gray tanager

| Model | QAIC | ΔQAIC | Model weight |
|--|-------------|--------------------------------|---------------------|
| $\Psi(\text{Forest}) p(\text{Forest})$ | 182.05 | 0.00 | 0.3944 |
| $\Psi(.) p(\text{Forest})$ | 182.64 | 0.59 | 0.2937 |
| $\Psi(\text{Forest}) p(.)$ | 183.29 | 1.24 | 0.2122 |
| $\Psi(.) p(.)$ | 184.80 | 2.75 | 0.0997 |

g. Boat-billed flycatcher

| Model | AIC | ΔAIC | Model weight |
|--|------------|-------------------------------|---------------------|
| $\Psi(.) p(.)$ | 153.70 | 0.00 | 0.5739 |
| $\Psi(\text{Forest}) p(.)$ | 155.16 | 1.46 | 0.2766 |
| $\Psi(.) p(\text{Forest})$ | 156.97 | 3.27 | 0.1119 |
| $\Psi(\text{Forest}) p(\text{Forest})$ | 159.15 | 5.45 | 0.0376 |

h. Golden-tailed sapphire

| Model | QAIC | ΔQAIC | Model weight |
|--|-------------|--------------------------------|---------------------|
| $\Psi(.) p(.)$ | 33.60 | 0.00 | 0.7099 |
| $\Psi(.) p(\text{Forest})$ | 36.91 | 3.31 | 0.1356 |
| $\Psi(\text{Forest}) p(.)$ | 36.93 | 3.33 | 0.1343 |
| $\Psi(\text{Forest}) p(\text{Forest})$ | 40.72 | 7.12 | 0.0202 |

i. Green and Gold tanager

| Model | AIC | ΔAIC | Model weight |
|--|------------|-------------------------------|---------------------|
| $\Psi(.) p(.)$ | 150.00 | 0.00 | 0.7529 |
| $\Psi(.) p(\text{Forest})$ | 153.30 | 3.30 | 0.1446 |
| $\Psi(\text{Forest}) p(.)$ | 154.00 | 4.00 | 0.1019 |
| $\Psi(\text{Forest}) p(\text{Forest})$ | 164.24 | 14.24 | 0.0006 |

j. Magpie tanager

| Model | AIC | ΔAIC | Model weight |
|--|------------|-------------------------------|---------------------|
| $\Psi(.) p(.)$ | 128.79 | 0.00 | 0.4932 |
| $\Psi(.) p(\text{Forest})$ | 130.03 | 1.24 | 0.2653 |
| $\Psi(\text{Forest}) p(\text{Forest})$ | 131.37 | 2.58 | 0.1358 |
| $\Psi(\text{Forest}) p(.)$ | 131.87 | 3.08 | 0.1057 |

k. Olivaceous woodcreeper

| Model | AIC | ΔAIC | Model weight |
|------------------------------------|------------|-------------------------------|---------------------|
| $\Psi(\cdot)$ p(\cdot) | 172.43 | 0.00 | 0.7284 |
| $\Psi(\text{Forest})$ p(\cdot) | 175.70 | 3.27 | 0.1420 |
| $\Psi(\cdot)$ p(Forest) | 176.23 | 3.80 | 0.1089 |
| $\Psi(\text{Forest})$ p(Forest) | 179.55 | 7.12 | 0.0207 |

l. Orange-bellied euphonia

| Model | QAIC | ΔQAIC | Model weight |
|------------------------------------|-------------|--------------------------------|---------------------|
| $\Psi(\cdot)$ p(\cdot) | 72.55 | (-) 0 | 0.6813 |
| $\Psi(\cdot)$ p(Forest) | 75.44 | 2.89 | 0.1606 |
| $\Psi(\text{Forest})$ p(\cdot) | 75.78 | 3.23 | 0.1355 |
| $\Psi(\text{Forest})$ p(Forest) | 79.36 | 6.81 | 0.0226 |

m. Palm tanager

| Model | AIC | ΔAIC | Model weight |
|------------------------------------|------------|-------------------------------|---------------------|
| $\Psi(\cdot)$ p(Forest) | 148.21 | 0.00 | 0.6253 |
| $\Psi(\cdot)$ p(\cdot) | 150.00 | 1.79 | 0.0255 |
| $\Psi(\text{Forest})$ p(Forest) | 152.21 | 4.00 | 0.0846 |
| $\Psi(\text{Forest})$ p(\cdot) | 154.00 | 5.79 | 0.0346 |

n. Paradise tanager

| Model | AIC | ΔAIC | Model weight |
|------------------------------------|------------|-------------------------------|---------------------|
| $\Psi(\cdot)$ p(Forest) | 208.29 | 0.00 | 0.8788 |
| $\Psi(\text{Forest})$ p(Forest) | 212.29 | 4.00 | 0.1189 |
| $\Psi(\cdot)$ p(\cdot) | 220.48 | 12.19 | 0.0020 |
| $\Psi(\text{Forest})$ p(\cdot) | 224.48 | 16.19 | 0.0003 |

o. Russet-backed oropendola

| Model | AIC | ΔAIC | Model weight |
|------------------------------------|------------|-------------------------------|---------------------|
| $\Psi(\cdot)$ p(Forest) | 268.25 | 0.00 | 0.5565 |
| $\Psi(\cdot)$ p(\cdot) | 269.33 | 1.08 | 0.3243 |
| $\Psi(\text{Forest})$ p(Forest) | 272.25 | 4.00 | 0.0753 |
| $\Psi(\text{Forest})$ p(\cdot) | 273.33 | 5.08 | 0.0439 |

p. Silver-beaked tanager

| Model | QAIC | ΔQAIC | Model weight |
|------------------------------------|-------------|--------------------------------|---------------------|
| $\Psi(\cdot)$ p(Forest) | 172.87 | 0.00 | 0.8795 |
| $\Psi(\text{Forest})$ p(Forest) | 176.87 | 4.00 | 0.1190 |
| $\Psi(\cdot)$ p(\cdot) | 186.73 | 13.86 | 0.0009 |
| $\Psi(\text{Forest})$ p(\cdot) | 187.30 | 14.43 | 0.0006 |

q. Violaceous jay

| Model | QAIC | ΔQAIC | Model weight |
|------------------------------------|-------------|--------------------------------|---------------------|
| $\Psi(\cdot)$ p(Forest) | 100.17 | 0.00 | 0.5929 |
| $\Psi(\cdot)$ p(\cdot) | 101.61 | 1.44 | 0.2886 |
| $\Psi(\text{Forest})$ p(Forest) | 104.17 | 4.00 | 0.0802 |
| $\Psi(\text{Forest})$ p(\cdot) | 105.65 | 5.48 | 0.0383 |

r. Yellow-rumped cacique

| Model | AIC | ΔAIC | Model weight |
|------------------------------------|------------|-------------------------------|---------------------|
| $\Psi(\text{Forest})$ p(\cdot) | 116.42 | 0.00 | 0.4554 |
| $\Psi(\text{Forest})$ p(Forest) | 116.72 | 0.30 | 0.3920 |
| $\Psi(\cdot)$ p(\cdot) | 119.50 | 3.08 | 0.0976 |
| $\Psi(\cdot)$ p(Forest) | 120.65 | 4.23 | 0.0549 |

Table S3. Proportion of three forests occupied for 18 bird species, calculated as model average estimates of $\Psi \pm$ unconditional standard error.

| Species | Manual restoration | Natural regrowth | Shifting cultivation |
|---------------------------|---------------------------|-------------------------|-----------------------------|
| Bananaquit | 0.998±0.023 | 1.000 | 0.998±0.023 |
| Bay-headed tanager | 0.968±0.091 | 0.997±0.009 | 0.997±0.009 |
| Black-faced dacnis | 0.842±0.235 | 0.893±0.173 | 0.851±0.229 |
| Black vulture | 0.319±0.305 | 0.439±0.380 | 0.319±0.305 |
| Blue-headed tanager | 1.000 | 1.000 | 1.000 |
| Blue-gray tanager | 0.975±0.0483 | 0.975±0.0483 | 0.617±0.317 |
| Boat-billed flycatcher | 0.966±0.093 | 0.966±0.093 | 0.860±0.206 |
| Golden-tailed sapphire | 0.280±0.150 | 0.218±0.148 | 0.249±0.138 |
| Green and Gold tanager | 0.9997±0.0006 | 1.000 | 1.000 |
| Magpie tanager | 0.909±0.170 | 0.914±0.155 | 0.834±0.210 |
| Olivaceous woodcreeper | 0.978±0.056 | 1.000 | 1.000 |
| Orange-bellied euphonia | 0.956±0.096 | 1.000 | 1.000 |
| Palm tanager | 1.000 | 1.000 | 1.000 |
| Paradise tanager | 1.00±0.0003 | 1.00±0.0003 | 1.00±0.0003 |
| Russett-backed oropendola | 1.000 | 1.000 | 0.974±0.052 |
| Silver-beaked tanager | 0.9999±0.0003 | 0.9999±0.0003 | 0.9996±0.0008 |
| Violaceous jay | 0.9991±0.0043 | 0.9991±0.0043 | 0.9991±0.0043 |
| Yellow-rumped cacique | 0.988±0.005 | 0.988±0.034 | 0.331±0.289 |

Table S4. Model averaged estimates (p) \pm unconditional standard error of detection probability of bird species by forest type.

| Species | Manual Restoration | Natural Regrowth | Shifting Cultivation |
|------------------------------|-------------------------------|-----------------------------|---------------------------------|
| Bananaquit | 0.107 \pm 0.026 | 0.108 \pm 0.026 | 0.124 \pm 0.035 |
| Bay-headed tanager | 0.073 \pm 0.035 | 0.127 \pm 0.039 | 0.279 \pm 0.068 |
| Black-faced dacnis | 0.072 \pm 0.087 | 0.114 \pm 0.087 | 0.325 \pm 0.090 |
| Black vulture | 0.404 \pm 0.521 | 0.404 \pm 0.528 | 0.510 \pm 0.074 |
| Blue-gray tanager | 0.447 \pm 0.0558 | 0.236 \pm 0.138 | 0.453 \pm 0.067 |
| Boat-billed flycatcher | 0.139 \pm 0.031 | 0.134 \pm 0.033 | 0.139 \pm 0.032 |
| Golden-tailed sapphire | 0.419 \pm 0.075 | 0.437 \pm 0.084 | 0.367 \pm 0.130 |
| Magpie tanager | 0.091 \pm 0.034 | 0.142 \pm 0.064 | 0.104 \pm 0.035 |
| Olivaceous woodcreeper | 0.152 \pm 0.028 | 0.155 \pm 0.029 | 0.152 \pm 0.029 |
| Orange-bellied euphonia | 0.097 \pm 0.027 | 0.105 \pm 0.025 | 0.111 \pm 0.030 |
| Palm tanager | 0.101 \pm 0.033 | 0.094 \pm 0.035 | 0.195 \pm 0.070 |
| Paradise tanager | 0.132 \pm 0.039 | 0.208 \pm 0.048 | 0.449 \pm 0.071 |
| Russett-backed oropendola | 0.464 \pm 0.069 | 0.367 \pm 0.056 | 0.354 \pm 0.067 |
| Silver-beaked tanager | 0.276 \pm 0.051 | 0.070 \pm 0.030 | 0.510 \pm 0.072 |
| Violaceous jay | 0.247 \pm 0.051 | 0.074 \pm 0.034 | 0.223 \pm 0.059 |
| Yellow-rumped cacique | 0.122 \pm 0.034 | 0.163 \pm 0.091 | 0.091 \pm 0.041 |